

The observed infall of galaxies towards the Virgo cluster

I. D. Karachentsev^{1*}, O. G. Nasonova (Kashibadze)^{1†}

¹*Special Astrophysical Observatory of the Russian Academy of Sciences, Nizhnij Arkhyz, KChR, 369167, Russia*

11 February 2010

ABSTRACT

We examine the velocity field of galaxies around the Virgo cluster induced by its overdensity. A sample of 1792 galaxies with distances from the Tip of the Red Giant Branch, the Cepheid luminosity, the SNIa luminosity, the surface brightness fluctuation method, and the Tully-Fisher relation has been used to study the *velocity-distance* relation in the Virgocentric coordinates. Attention was paid to some observational biases affected the Hubble flow around Virgo.

We estimate the radius of the zero-velocity surface for the Virgo cluster to be within $(5.0 - 7.5)$ Mpc corresponding to $(17 - 26)^\circ$ at the mean cluster distance of 17.0 Mpc. In the case of spherical symmetry with cosmological parameter $\Omega_m = 0.24$ and the age of the Universe $T_0 = 13.7 Gyr$, it yields the total mass of the Virgo cluster to be within $M_T = (2.7 - 8.9) \cdot 10^{14} M_\odot$ in reasonable agreement with the existing virial mass estimates for the cluster.

Key words: galaxies: clusters: individual: Virgo – dark matter.

1 INTRODUCTION

The gravitational action of mass of a solitary system of galaxies leads to deceleration of the local Hubble flow. As a result, the line of average velocity of neighboring galaxies relative to the center of a cluster (or a group) deviates from the linear Hubble relation, going to negative values at small distances $R < R_0$. Here, R_0 means the radius of the zero-velocity surface, which separates the galaxy system against the global cosmic expansion. As it was shown by Lynden-Bell (1981) and Sandage (1986), in the simplest case of spherical symmetry with

* E-mail:ikar@sao.ru

† E-mail:phiruzi@gmail.com

cosmological parameter $\Lambda = 0$ the radius R_0 depends only on the total mass of a group M_T and the age of the Universe T_0 :

$$M_T = (\pi^2/8G) \cdot R_0^3 \cdot T_0^{-2}, \quad (1)$$

where G is the gravitational constant. Measuring R_0 via distances and radial velocities of galaxies outside the virial radius of the system R_{vir} , one can determine the total mass of the system independent of its virial mass estimate. Note that both the methods deriving mass from internal and from external galaxy motions correspond to different linear scales where R_0 is roughly 4 times as large as the virial radius. In reality, galaxy groups and clusters do not have perfect spherical symmetry, and cosmology with $\Lambda = 0$ is not true.

Numerous measurements of distances to nearby galaxies obtained recently with the Hubble Space Telescope (HST) allowed us to investigate the Hubble flow around the Local Group and other proximate groups (Karachentsev et al. 2002, Karachentsev & Kashibadze 2006, Karachentsev et al. 2006, 2009). The radii R_0 obtained from observations for nearby groups around: the Milky Way & Andromeda (the Local Group), M81, Centaurus A, Maffei & IC 342, NGC 253 (Sculptor filament), and NGC 4736 (Canes Venatici I cloud) are ranged within (0.7–1.4 Mpc). The average ratio of total-to-virial masses for these six groups, derived from R_0 via eq. (1) and from R_{vir} , turns out to be $\langle M_T/M_{vir} \rangle = 0.60 \pm 0.15$ (Karachentsev, 2005). But as it was noticed by Peirani & Pacheco (2006, 2008) and Karachentsev et al. (2007), in a flat universe dominated by dark energy the resulting $M_T(R_0)$ mass is higher than that derived from the canonical Lemaître-Tolman eq. (1). In the "concordant" cosmological model with Λ -term and Ω_m as a matter component eq. (1) takes a form

$$M_T = (\pi^2/8G) \cdot R_0^3 \cdot H_0^2 / f^2(\Omega_m), \quad (2)$$

where

$$f(\Omega_m) = (1 - \Omega_m)^{-1} - (\Omega_m/2) \cdot (1 - \Omega_m)^{-3/2} \cdot \operatorname{arccosh}[(2/\Omega_m) - 1]. \quad (3)$$

Assuming $\Omega_m = 0.24$ and $H_0 = 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$ that corresponds to $T_0 = 13.7 \text{ Gyr}$ (Spergel et al. 2007), one can rewrite (2) as

$$(M_T/M_\odot)_{0.24} = 2.12 \cdot 10^{12} (R_0/\text{Mpc})^3. \quad (4)$$

It yields the mass that is 1.5 as large as derived from the classic eq. (1). This correction

leads to a good agreement in average between the R_0 mass estimates and virial masses for the abovementioned galaxy groups.

For galaxies around the nearest cluster in Virgo, expected velocity deviations from the pure Hubble flow (so-called Virgocentric infall) were regarded in dynamical models by Hoffman et al. (1980), Tonry & Davis (1981) and Hoffman & Salpeter (1982). These authors note that with the virial mass of the Virgo cluster $M_{vir} \sim 6 \cdot 10^{14} M_{\odot}$, the radius of the zero-velocity surface around the cluster amounts to $\sim 27^\circ$, i.e. the infall zone covers nearly 1 steradian of the sky. According to Hoffman et al. (1980), the observed decrease of radial velocity dispersion within the angular distance $\Theta = [0 - 24]^\circ$ from the Virgo center for 228 galaxies agrees, in the main, with the Virgocentric infall pattern for the cluster mass mentioned above. Tully & Shaya (1984) considered the phenomena of infall of galaxies towards Virgo both in the point-mass and distributed-mass models for the cluster with different values of the cosmological parameter Ω_{Λ} and the age of the Universe T_0 . Using Tully-Fisher distance estimates for 19 galaxies inside the virial radius of $\sim 6^\circ$ and 14 galaxies outside it, the authors ascertain the expected infall with $R_0 \sim 28^\circ$.

Later, Tonry et al. (2000, 2001) developed a model of the Virgocentric flow basing on accurate distance measurements for 300 E and S0 galaxies from surface brightness fluctuations. Their model fits well the observational data on galaxy distances and radial velocities for the Virgo cluster distance of 17.0 Mpc and its total mass of $7 \cdot 10^{14} M_{\odot}$. According to their model, our Local Group has a peculiar velocity of 139 km s^{-1} directed towards the Virgo center.

Teerikorpi et al. (1992), Ekholm et al. (1999) and Ekholm et al. (2000) examined the Virgocentric flow with different models of density distribution in the cluster and inferred expected relations between velocities and distances of galaxies relative to the cluster center. Using Cepheid distances to 23 galaxies and Tully-Fisher distances to 96 galaxies, the authors conclude that the radius of the zero-velocity surface ranges from 20° to 31° , and the total cluster mass is equal to $(1 \div 2)$ of its virial value.

During the last decade, the observational database on distances to galaxies in a wide vicinity of the Virgo cluster has grown significantly, allowing to determine R_0 and, therefore, the total mass of the Virgo cluster with better accuracy.

2 THE OBSERVATIONAL SAMPLES

To examine phenomena of the Virgocentric flow, we used distance moduli of galaxies from different publications preferring more precise measurements. The main data sources are listed below.

A) Taking luminosity of the Tip of the Red Giant Branch (TRGB) as a standard is the most efficient and the most universal method to determine distances to nearby galaxies, as it is practically independent of their morphological type. Being applied to galaxy images in two or more photometric bands obtained with WFPC2 or ACS cameras at the HST, the TRGB method yields an accuracy of distance measurements of $\sim 7\%$, as it was founded by Rizzi et al. (2007). A consolidated list of distances for the Local Volume galaxies is presented in the Catalog of Neighboring Galaxies (= CNG, Karachentsev et al. 2004). The CNG sample of 451 galaxies has been collected based on two conditions: galaxy distance $D < 10$ Mpc, if a galaxy has individual distance estimate; otherwise galaxy radial velocity with respect to the Local Group $V_{LG} < 550$ km s $^{-1}$. Below we use from CNG the galaxies with only TRGB or Cepheid distances, supplying them with new TRGB distances from recent publications (Karachentsev et al. 2006, Tully et al. 2006).

B) The surface brightness fluctuation method, SBF, applying to early type galaxies, assumes that the old stellar population (RGB) is prevailing in a total luminosity, and the galaxy structure does not suffer from irregularities due to dust clouds. Using this approach, Tonry et al. (2001) determined SBF distances to 300 E and S0 galaxies with a typical errors of $\sim 12\%$. This sample is distributed over the whole sky extending to $cz \sim 4000$ km s $^{-1}$ with a median velocity of 1800 km s $^{-1}$.

C) Mei et al.(2007) undertook a two-color ACS/HST imaging survey for 100 early-type galaxies situated in the Virgo cluster core (the ACSVCS project). They derived precise SBF distances to 84 E, S0 galaxies with a typical error of 8%, and revealed a 3D-shape of the Virgo cluster to be a slightly triaxial ellipsoid with axis ratios of (1:0.7:0.5). We expanded the ACSVCS sample with other precise SBF and TRGB distance measurements in the Virgo core made by Neilsen & Tsvetanov (2000) and Caldwell (2006) with ACS/HST and by Jerjen et al. (2004) with the VLT. That yields us the total "ACSVCS+" sample of 116 galaxies.

D) In wide vicinity of the Virgo cluster there are 22 galaxies with distances measured by Tonry et al. (2003) via SNIa. This sample is small but has the distance error of only 5%.

E) Based on the "2MASS Selected Flat Galaxy Catalog" (Mitronova et al. 2004 =

2MFGC), Kashibadze (2008) determined distances to 402 spiral edge-on galaxies with radial velocities $< 3000 \text{ km s}^{-1}$. A multiparametric NIR Tully-Fisher relation was applied to them, yielding a typical distance error of $\sim 20\%$. The zero point of the luminosity - line width relation was calibrated by 15 galaxies with Cepheid and TRGB distance measures.

F) Finally, the former samples of galaxies were supplemented with a compilation of distances by Tully et al. (2008, 2009) that have been obtained from optical (B, R or I band) one-parametric Tully-Fisher relations. This compilation relies on numerous HI-line and photometric observations carried out by Methewson & Ford (1996), Haynes et al. (1999), Tully & Pierce (2000), Koribalski et al. (2004), Springob et al. (2005), Theureau et al. (2006) and other authors. Zero points of the data were recalibrated by a set of 40 galaxies with known Cepheid and TRGB distances. As a last step, we used also distances from a very big and important SFI++ sample (Springob et al. 2007), which have not been included in Tully et al. (2009) compilation. At total, we used distance estimates for 941 spiral galaxies whose radial velocities were limited by 3000 km s^{-1} . The typical distance error for them is $\sim 20\%$, though there are some cases with much higher errors due to uncertainties of galaxy inclination, presence of interacting companions or HI profiles of low quality.

A substantial overlap between the two last *luminosity-linewidth* samples provides confirmation that their zero points are the same and gives rms agreement per measure of 0.40 mag. As it is seen from Fig. 1 by Tully et al. (2008), there is an excellent agreement in distance moduli between the *luminosity-linewidth* and other (Cepheid, TRGB, SBF and SNIa) measures. In particular, for 12 galaxies in our list with both TF and TRGB or SNIa moduli the mean distance difference is $(-1.4 \pm 1.2) \text{ Mpc}$, while for 20 galaxies with SBF or TRGB moduli the average difference is only $(0.2 \pm 0.2) \text{ Mpc}$.

Following the previous authors (Tully & Shaya 1984, Ekholm et al. 2000) we have formed a composite sample of galaxies limiting their angular separation from the Virgo center to $\Theta < 30^\circ$. We have considered the radiogalaxy Virgo A = NGC 4486 to be the physical center of the cluster as its position is close to the center of X-ray emitting gas. The total number of galaxies in this cone volume with apex angle $\Theta < 30^\circ$ is 630 (the *2D sample*).

Limiting the angular separation of galaxies from the Virgo center introduces some selection effects into the Virgocentric flow analysis. For this reason we have used also another way to form the observational sample, considering galaxies with spatial distances from the Virgo cluster center $R_{vc} < 30 \text{ Mpc}$ (the *3D sample*). This approach suffers a drawback too because galaxy distances are measured with errors and their significance is different at the

proximate and the distant boundary of the spherical volume (the so-called *Malmquist bias*). The total number of galaxies in our 3D sample amounts to 1792, and the fractions of diverse subsamples differ significantly from that in the 2D sample.

Table 1 presents the summary of observational data that we have used. The first column indicates subsample kinds, the second one gives typical distance errors expressed in magnitudes. Columns (3) and (5) contain numbers of galaxies in the cone (2D) or in the spherical (3D) volume. The *sample goodness* G , defined as $G = (N/100)^{1/2} \cdot \sigma_m^{-1}$, is a useful parameter which characterizes a statistical weight of a certain sample (Kudrya et al. 2003). Goodness values are indicated in columns (4) and (6). For example, the subsample of galaxies with SNIa distances is scanty but its statistical significance is comparable with that of other samples because of higher accuracy of distance measurements. As one can see, the galaxy subsample ACSVCS+ has the maximum statistical weight in the 2D set, however almost all these galaxies are concentrated within the virial radius. In the 3D set the highest goodness corresponds to the TRGB sample, but its majority is crowded on the nearby side of the examined volume. The last two TF samples exhibit a significant increase in number going from the 2D to the 3D samples that is caused by the well-known effect of morphological segregation of late-type vs. early-type galaxies along the cluster radius.

3 RADIUS OF THE ZERO-VELOCITY SURFACE R_0

The virial radius of the Virgo cluster $R_{vir} = 1.8$ Mpc (Hoffman et al. 1980) corresponds to its angular scale of 6.0° , assuming the average distance to the cluster members to be 17.0 Mpc. Radial velocities and distances relative to the Local Group (LG) centroid for 259 galaxies in this zone are represented in the top panel of Fig. 1. Here, precise distances for most the galaxies were obtained within the special survey ACSVCS with HST (Mei et al. 2007). The Virgo cluster members, located in the distance range from 14 to 20 Mpc, demonstrate a radial velocity scatter from -800 up to $+2300$ km s $^{-1}$. Foreground galaxies are scarcely presented on the panel while background objects tend to lie below the linear Hubble regression with the global Hubble parameter $H_0 = 72$ km s $^{-1}$ Mpc $^{-1}$ (Spergel et al. 2007), showing thereby the expected effect of infall into the Virgo cluster from the opposite side. The centroid of galaxies forming the "virial column" at $[17.0 \pm 1.8]$ Mpc, marked by vertical lines, has a mean velocity $+1004 \pm 70$ km s $^{-1}$ versus the expected Hubble velocity of $+1224$ km s $^{-1}$ at the distance of 17.0 Mpc, which can be explained by a peculiar motion

of the Local Group $\sim 220 \pm 70 \text{ km s}^{-1}$ directed towards Virgo. The dotted and solid S-shaped curves correspond to a Hubble flow perturbed by a point-like mass of $2.7 \cdot 10^{14} M_{\odot}$ and $8.9 \cdot 10^{14} M_{\odot}$ (as the limiting cases discussed below) for the line-of-sight passing exactly through the cluster center.

The distributions of radial velocities and distances for remaining galaxies of the 2D sample in close surroundings of Virgo, ($6^{\circ} < \Theta < 15^{\circ}$) and in a distant periphery ($15^{\circ} < \Theta < 30^{\circ}$), are shown in the middle and the bottom panels of Fig. 1. Here, the solid and dotted S-shaped lines having lower amplitudes describe the behavior of perturbed Hubble flow at angular distances Θ equal 6° and 15° , respectively. These panels display some signs of the infall effect too, however, in front of the Virgo, the expected infall is seen barely.

Considering a set of such kind Hubble diagrams with their different amplitudes of S-shaped waves decreasing with the angular distance Θ , one can find the quantity of the cluster mass that fits the observed infall pattern in the best way. But this approach seems us to be not transparent enough. In order to determine R_0 and the total cluster mass via it, we have converted our observational data into distances and velocities expressed relative to the cluster center. The top panel of Fig. 2 shows the layout of a galaxy (G) relative to the observer (LG) and the cluster center (C) with angular separation Θ from the cluster center. The spatial distance of the galaxy from the center therefore is

$$R_{vc}^2 = R_g^2 + R_c^2 - 2R_g R_c \cos \Theta. \quad (5)$$

Assuming that the galaxy and the cluster center are involved in an almost unperturbed Hubble flow (Hf case) with negligible peculiar velocities, we can state the mutual velocity difference between G and C in projection onto the stright line connecting them as

$$V_{vc} = V_g \cos \lambda - V_c \cos \mu, \quad (6)$$

where $\mu = \lambda + \Theta$ and $\tan \lambda = R_c \sin \Theta / (R_g - R_c \cos \Theta)$.

The distribution of galaxies in the Virgocentric reference frame $\{V_{vc}, R_{vc}\}$ is represented in Fig. 3. The only 391 galaxies with angles λ obeying $\lambda < 45^{\circ}$ or $\lambda > 135^{\circ}$ from the whole 2D sample are shown here. Selecting galaxies situated approximately in front and behind the cluster is meant to reduce the role of tangential velocity componens which remain still unknown. The polygon curve traces the running median with a window of 1 Mpc. The median follows roughly the linear Hubble regression with $H_0 = 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (the

inclined dashed line) at middle Virgocentric distances of ~ 15 Mpc, but tends to deviate from the $H_0 \cdot R_{vc}$ line at smaller scales, crossing the zero-velocity line at $R_0 \simeq 6$ Mpc. The behaviour of the running median at large scales is strongly skewed by a selection effect due to the adopted limit for galaxy velocities $V_{LG} < 3000 \text{ km s}^{-1}$.

Another approach can be used also for converting the observational radial velocity of a galaxy V_g into its Virgocentric velocity (the case of pure Virgocentric flow, V_f). If the galaxy is not involved in the general cosmological expansion but is falling instead towards the Virgo cluster with a velocity V_{in} (Fig. 2b), than its radial velocity relative to the observer will be expressed as

$$V_g = V_c \cdot \cos \Theta - V_{in} \cos \lambda, \quad (7)$$

and the infall velocity V_{in} itself can be written as

$$V_{in} = (V_c \cos \Theta - V_g) \sec \lambda, \quad (8)$$

where the angles Θ and λ are shown in the Fig. 2b. Evidently, the discrepancy between these two extreme approaches, Hf and Vf , decreases when λ tends to 0 or to 180° .

As can be easily seen, selecting galaxies both in the cone with apex angle $\Theta = 30^\circ$ and the angle λ entails a loss of galaxies at large Virgocentric distances which becomes apparent in the top right corner of the Fig. 3. A selection of galaxies in the volume $R_{vc} < 30$ Mpc (3D sample) reduces the bias appreciably. Fig. 4 represents the sky distribution of galaxies with known Virgocentric distances up to 30 Mpc in equatorial coordinates. The galaxies of this 3D sample are marked as circles and their diameters indicate three distance ranges: (0–12) Mpc, (12–22) Mpc, and more than 22 Mpc from the observer. This map exhibits that the selection of galaxies by their angle $\Theta < 30^\circ$ brings some systematic skews dependent on distance. In particular foreground Virgo galaxies are preferentially removed by this angular selection.

The Hubble diagram for 1792 galaxies of the 3D sample is represented in Fig. 5. The symbols for objects from different sources of distance data are the same as for Fig. 1. The number of galaxies in the 3D sample is roughly 3 times as large as in the 2D sample. It is noteworthy, they populate the crucial regions in front and behind the Virgo cluster more thoroughly giving us an opportunity for more detailed analyses of the Virgocentric infall. The velocity - distance relation for these galaxies with respect to the cluster center is shown

in Fig. 6. Its top panel corresponds to the assumption of pure Hubble flow of galaxies (*Hf* case) while the bottom panel represents the case of pure radial motions towards the Virgo center (*Vf*). As previously, the galaxies located far away from the line of sight crossing the cluster center, i.e. with $45^\circ < \lambda < 135^\circ$, are eliminated in order to reduce the role of unknown tangential velocity components. This condition diminishes the number of sampled galaxies by 42%.

Comparison the bottom panel of Fig. 6 with the top one shows that switching from the *Hf* case to the *Vf* case does not lead to any dramatic changes in the Hubble flow pattern given in the Virgocentric coordinates. Some galaxies move along the vertical axis appreciably but the total behaviour of the running medians traces the infall of galaxies towards the cluster in a similar way. The asymptotic tendency of the median at large distances R_{vc} looks much more regular for the 3D sample than for the 2D one.

Our elimination of galaxies with $45^\circ < \lambda < 135^\circ$ is a little arbitrary. To estimate the response of the Virgocentric flow pattern to changing this condition we have imposed also a more rigid constraint, eliminating galaxies with $30^\circ < \lambda < 150^\circ$. The corresponding diagrams for *Hf* and *Vf* cases are represented in two panels of Fig. 7. As it is seen, the more severe selection of galaxies by λ reduces their number by more than a half. However, the behaviour of the running median is almost the same.

In the bottom panel of Fig. 7 every galaxy from the samples A,B,C, and D (see Table 1) having a distance error within 12% is supplied by error bars indicating where the galaxy should be situated if its distance from the observer changes by $\pm 1\sigma_D$. As expected, these error bars are much longer for galaxies situated behind the cluster. Changing a galaxy distance by $\pm 12\%$ leads in some regions of the $\{V_{vc}, R_{vc}\}$ diagram to significant displacement of the galaxy along the both Virgocentric coordinates, and therefore to appreciable galaxy skips relative to the zero-velocity line. (We do not include the longer error bars for the Tully-Fisher distances).

To quantify the uncertainties on the running median curves plotted in Figs. 6 and 7, we generated extensive sets of bootstrap realisations. Their results permit us to estimate the radius of the zero velocity surface and also its rms error presented in Table 2. The table columns contain: (1) size of the window for a running median, taken to be 0.8, 1.0, and 1.2 Mpc; (2,3) the mean radius of the zero velocity surface R_0 and its rms scatter obtained with regarding to the assumptions on the pure Hubble flow (*Hf*) and the pure Virgocentric flow (*Vf*), respectively; here only four samples: A,B,C, and D with precise distance moduli were

taken into account; (4,5) the same quantities for the samples E and F with Tully-Fisher distances; (6,7) the radii R_0 and their errors for the total set of available data on galaxy distances. We shall discuss the interpretation of these different estimates of R_0 in the next section.

4 DISCUSSION AND CONCLUSIONS

We have analysed available observational data on distances and radial velocities of galaxies in wide surroundings of the Virgo cluster in order to study the Virgocentric infall. The main purpose of the present paper is to determine the radius of the zero-velocity surface R_0 that separates the cluster against the global cosmic expansion. By using this observational quantity we are able to estimate the total mass of the Virgo concentrated within R_0 and compare it with the virial mass estimates corresponding to roughly 4 times lower scale, R_{vir} . Based on the Hubble diagrams transformed into Virgocentric coordinates and the results of our bootstrap numerical experiments given in Table 2, we derive R_0 to be in the range of 5.0 – 7.5 Mpc with a typical random error of 1.0 Mpc. The derived value of R_0 can be affected by some systematic circumstances.

Analysis of the Hubble diagrams in the Virgocentric coordinates suffers the lack of data on tangential velocity components of the galaxies. We tried to overcome this drawback with assumptions regarding a dominant type of galaxy motions in the Virgo proximity. Conversion of the observed radial velocities of galaxies into their Virgocentric velocities was performed by us under two extreme kinematic assumptions: almost unperturbed Hubble flow (Hf) or almost pure radial flow towards Virgo (Vf). We found that adopting one or another scheme doesn't change significantly a general pattern of the Virgocentric infall. Calculating velocities in the Vf case yields, at the average, some smaller values of V_{vc} , which causes a slightly larger (+0.7 Mpc) value of R_0 .

The R_0 quantities presented in Table 2 tend to be correlated with the smoothing window size. When the window changes from 0.8 Mpc to 1.2 Mpc, the radius decreases on 0.3 Mpc at average. Variation the window size in wider range: from 0.5 Mpc to 2.0 Mpc still leaves R_0 within its random errors.

The most noticeable systematic variations in the radius R_0 are seen in dependence on galaxy samples. The use of samples of galaxies with only precise distance measurements (via Cepheids, TRGB, SBF, and SNIa) yields the radius R_0 within 6.5 – 7.5 Mpc, while the use of

less precise Tully-Fisher distances leads to R_0 estimates around 5.2 Mpc. The physical origin of this difference is clear. Probably, the instance is just related with the fact that a typical distance error for Tully-Fisher method ($\sim 20\%$) corresponds at the Virgo distance to a linear scale of 3.4 Mpc that comparable with the virial diameter of the cluster (3.6 Mpc). Large random errors can throw galaxies over the "virial column" diluting the shape of S-wave infall. Nevertheless, we should stress that two completely independent sets of observational data on distances to early-type and late-type galaxies lead to quite compatible values of R_0 . As can be seen from Table 1, two samples with Tully-Fisher distances (E+F) are about 2 times as large in galaxy number as the samples (A+B+C+D) with precise distances. Because we estimate R_0 based on the running median without regard to galaxy weights (distance errors), the radius R_0 for the total sample turns out to be closer to that for the former samples.

Assuming R_0 for the Virgo cluster in the range of (5.0 – 7.5) Mpc, we obtain from eq. (4) the total mass of the cluster to be within $M_T = (2.7 - 8.9) \cdot 10^{14} M_\odot$. This value is consistent with the virial mass estimates for the Virgo: $(5.5 \pm 1.4) \cdot 10^{14} M_\odot$ (Hoffman et al. 1980), $(7.5 \pm 1.5) \cdot 10^{14} M_\odot$ (Tully & Shaya 1984) and $7.0 \cdot 10^{14} M_\odot$ (Tonry et al. 2000) normalized to the same Hubble parameter $H_0 = 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$. It should be reminded that this agreement is on very different scales as the zero velocity radius, R_0 , is roughly 4 times as large as the virial radius. This means our results are consistent with there being **no additional mass** outside the virial radius of the Virgo cluster.

Two items are worth mentioning in conclusions. The stated method of estimating R_0 value from observational data relies on the assumptions that a cluster has spherically symmetric shape and that galaxy motions around the cluster are regular without a significant tangential component. Deviations from the simple spherical infall scenario have been considered both theoretically and observationally by many authors (Peebles 1980, Davis & Peebles 1983, Lilje et al. 1986, Tonry et al. 2000, Mould et al. 2000, Watkins et al. 2009). In particular, Tonry et al. (2000) studied a peculiar velocity field around the Local Group and Virgo cluster in the presence of second attractor (Great Attractor) situated in Hydra-Centaurus. As seen from their Figs. 20,21, the Great Attractor with a total mass of $9 \cdot 10^{15} M_\odot$ situated at a distance of 43 Mpc generates significant distortions in the velocity field over the area of ~ 1 steradian. Besides, Tully (1988) found so called effect of "Local Velocity Anomaly" arising by a push of the Local Void. According to Tully et al. (2009), a bulk motion of $\sim 260 \text{ km s}^{-1}$ toward the Supergalactic pole may be attributed to the Local Void. Recently, cosmic

velocity flows in the Local Universe were also studied by Erdogdu et al. (2006), Haugbolle et al. (2007) and Lavaux (2009).

We would emphasize that observational abilities to determine R_0 and M_T with a better accuracy have not been exhausted yet. With the derived R_0 around 7 Mpc, the zero-velocity surface is situated at a distance of about 10 Mpc from the observer, i.e. on the Local volume edge. Hence, one can expect to find some number of the LV galaxies with radial velocities $\sim (1000 - 2000)$ km s $^{-1}$, residing in front of Virgo. A new updated version of CNG (Karachentsev et al. 2010) contains about 40 candidate dwarf galaxies, like VCC1675, with appropriate radial velocities and rough distance estimates. Their precise distances could be easily measured via TRGB with facilities of ACS at HST in the one-object- per-one-orbit mode. The addition of more galaxies with measured distances in the range 9 – 11 Mpc from the Local Group would allow a better measurement of R_0 , in particular these galaxies could be used to better rule out the case where there is a large amount of mass outside the virial radius.

Acknowledgements. This work was supported by RFBR 07-02-00005, RFBR-DFG 06-02-04017 and CNRS grants. Authors thank Dmitry Makarov, Helene Courtois and Stefan Gottloeber for useful discussions. We would also like to thank the anonymous referees for their valuable comments which led to significant improvements in the paper.

REFERENCES

- Caldwell N., 2006, ApJ, 651, 822
 Davis M., Peebles P. J. E., 1983, ARA&A, 21, 109
 Ekholm T., Lanoix P., Teerikorpi P., Fouque P., Paturel G., 2000, A&A, 355, 835
 Ekholm T., Lanoix P., Teerikorpi P., Paturel G., Fouque P., 1999, A&A, 351, 827
 Erdogdu P., Lahav O., Huchra J. P. et al. 2006, MNRAS, 373, 45
 Haugbolle T., Hannestad S., Thomsen B. et al. 2007, ApJ, 661, 650
 Haynes M.P., Giovanelli R., Salzer J. J. et al. 1999, AJ, 117, 1668
 Hoffman G. L., Salpeter E. E., 1982, ApJ, 263, 485
 Hoffman G. L., Olson D. W., Salpeter E. E., 1980, ApJ, 242, 861
 Jerjen H., Binggeli B., Barazza F. D., 2004, AJ, 127, 771
 Karachentsev I. D., et al. 2010 (in preparation)
 Karachentsev I. D., Kashibadze O. G., Makarov D. I., Tully R. B., 2009, MNRAS, 393, 1265
 Karachentsev I. D., Kashibadze O. G., 2006, Astrophysics, 49, 3
 Karachentsev I. D., Dolphin A. E., Tully R. B. et al., 2006, AJ, 131, 1361
 Karachentsev I. D., 2005, AJ, 129, 178
 Karachentsev I. D., Karachentseva V. E., Huchtmeier W. K., Makarov D. I., 2004, AJ, 127, 2031 (CNG)
 Karachentsev I. D., Sharina M. E., Makarov D. I. et al., 2002, A&A, 389, 812



- Kashibadze O. G., 2008, *Astrophysics*, 51, 409
- Koribalski B. S., Staveley-Smith L., Kilborn V. A. et al. 2004, *AJ*, 128, 16
- Kudrya Yu. N., Karachentseva V. E., Karachentsev I. D., Mitronova S. N., et al., 2003, *A&A*, 407, 889
- Lavaux G., 2009, *asro-ph/0912.0516*
- Lilje P. B., Yahil A., Jones B. J., 1986, *ApJ*, 307, 91
- Lynden-Bell D., 1981, *The Observatory*, 101, 111
- Methewson D. S., Ford V. L., 1996, *ApJS*, 107, 97
- Mei S., Blakeslee J. P., Côte P., Tonry J. L., et al., 2007, *ApJ*, 655, 144
- Mitronova S. N., Karachentsev I. D., Karachentseva V. E., Jarrett T. H., Kudrya Yu. N., 2004, *Bull. Spec. Astrophys. Obs.*, 57, 5 (2MFGC)
- Mould J. R., Huchra J. P., Freedman W. L., et al., 2000, *ApJ*, 529, 786
- Neilsen E. H., Tsvetanov Z. I., 2000, *ApJ*, 536, 255
- Peirani S., Pacheco J. A., 2008, *A & A*, 488, 845
- Peirani S., Pacheco J. A., 2006, *New Astr.*, 11, 325
- Peebles P. J. E., 1980, *The Large Scale Structure of the Universe*, Princeton, Princeton University Press
- Rizzi L., Tully R. B., Makarov D. I., et al., 2007, *ApJ*, 661, 813
- Sandage A., 1986, *ApJ*, 307, 1
- Spergel D. N., et al., 2007, *ApJS*, 170, 377
- Springob C. M., Masters K. L., Haynes M. P. et al. 2007, *ApJS*, 172, 599
- Springob C. M., Haynes M. P., Giovanelli R., Kent B. R., 2005, *ApJS*, 160, 149
- Teerikorpi P., Bottinelli L., Gouguenheim L., Paturel G., 1992, *A&A*, 260, 17
- Theureau G., Martin J. M., Cognard L., et al. 2006, in *ASP conference ser.*, Vol. 351, ed. G. Gabriel, p. 429
- Tonry J. L., Schmidt B. P., Barris B., et al., 2003, *ApJ*, 594, 1
- Tonry J. L., Dressler A., Blakeslee J. P., et al., 2001, *ApJ*, 546, 681
- Tonry J. L., Blakeslee J. P., Ajhar E. A., Dressler A., 2000, *ApJ*, 530, 625
- Tonry J. L., Davis M., 1981, *ApJ*, 246, 680
- Tully R. B., Rizzi L., Shaya E. J., Courtois H. M., Makarov D. I., Jacobs B. A., 2009, *AJ*, 138, 323
- Tully R. B., Shaya E. J., Karachentsev I. D., Courtois H. M., et al., 2008, *ApJ*, 676, 184
- Tully R. B., Rizzi L., Dolphin A. E., et al., 2006, *AJ*, 132, 729
- Tully R. B., Pierce M. J., 2000, *ApJ*, 533, 744
- Tully R. B., 1988, *Nature*, 334, 209
- Tully R. B., Shaya E. J., 1984, *ApJ*, 281, 31
- Watkins R., Feldman H. A., Hudson M. J., 2009, *MNRAS*, 392, 743

Table 1. Distance datasets for galaxies in/around Virgo.

Sample	σ_m	N(2D)	G(2D)	N(3D)	G(3D)
A: TRGB + Ceph	0.15	37	4.1	264	10.8
B: SBF (Tonry)	0.25	81	3.6	189	5.5
C: ACSVCS+	0.17	113	6.2	113	6.3
D: SNIa (Tonry)	0.10	13	3.6	22	4.7
E: TF (IR)	0.40	56	1.9	260	4.0
F: TF (opt)	0.40	330	4.5	941	7.7

Table 2. Radius R_0 and its rms error (in Mpc) obtained from bootstrap realisations.

Window (Mpc)	Primary distances (samples A+B+C+D)		TF distances (samples E+F)		All samples	
	Hf	Vf	Hf	Vf	Hf	Vf
0.8	6.79	7.78	5.07	5.58	5.15	5.82
	± 1.12	± 1.13	± 1.04	± 1.13	± 0.96	± 1.04
1.0	6.59	7.54	4.87	5.38	5.00	5.65
	± 1.05	± 1.03	± 0.88	± 0.97	± 1.00	± 1.07
1.2	6.46	7.42	4.71	5.24	4.80	5.52
	± 1.01	± 0.97	± 0.78	± 0.87	± 0.95	± 1.03






Figure 1. The radial velocity vs. distance relation for galaxies in the Virgo cluster region with respect to the Local Group centroid. Galaxy samples with distances derived by different methods are marked by different symbols. The inclined line traces Hubble relation with the global Hubble parameter $H_0 = 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$. The vertical dashed lines outline the virial zone. Two S-shaped lines correspond to a Hubble flow perturbed by virial masses of $2.7 \cdot 10^{14} M_\odot$ (dotted) and $8.9 \cdot 10^{14} M_\odot$ (solid) as the limiting cases within confidence range. *Top:* the cluster core within angular distance $\Theta < 6^\circ$, both the S-shaped lines indicate the expected infall at $\Theta = 0^\circ$. *Middle:* wider surroundings with $6^\circ < \Theta < 15^\circ$, the S-shaped lines indicate the infall at $\Theta = 6^\circ$. The typical distance error bars for each dataset are shown. *Bottom:* peripheric regions with $15^\circ < \Theta < 30^\circ$, the S-shaped lines indicate the infall at $\Theta = 15^\circ$.




Figure 2. Galaxy (G) motion with respect to the cluster center (C) in the Local Group (LG) rest frame. *Top:* in the case of almost pure Hubble flow. *Bottom:* in the case of almost pure Virgo-centric infall.




Figure 3. Hubble flow around Virgo in the Virgocentric coordinates for 391 galaxies with $\Theta < 30^\circ$ and small angles λ regarding to the line of sight. The inclined dashed line corresponds to $H_0 = 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$, and the polygon curve traces the running median with a window of 1 Mpc. The dotted and solid curves correspond to a Hubble flow perturbed by a point-like mass of $2.7 \cdot 10^{14} M_\odot$ and $8.9 \cdot 10^{14} M_\odot$ respectively as the limiting cases.




Figure 4. Sky distribution of 1792 galaxies with Virgocentric distances $R_{vc} < 30$ Mpc in equatorial coordinates. Circles of different diameters indicate three distance ranges from the LG.




Figure 5. The Hubble diagram for 1792 galaxies with Virgocentric distances less than 30 Mpc. Symbol indications the same as in Fig. 1.




Figure 6. The Hubble flow in the Virgo-centric coordinates for 1045 galaxies with $R_{vc} < 30$ Mpc. *Top:* the case of almost pure Hubble flow. *Bottom:* the case of almost pure Virgo-centric infall. Only the galaxies with $\lambda < 45^\circ$ or $\lambda > 135^\circ$ are presented. The dotted and solid curves correspond to a Hubble flow perturbed by a point-like mass of $2.7 \cdot 10^{14} M_{\odot}$ and $8.9 \cdot 10^{14} M_{\odot}$ respectively as the limiting cases.




Figure 7. The same Hubble diagram as in Fig. 6, but only for 467 galaxies with the angles $\lambda < 30^\circ$ or $\lambda > 150^\circ$. In the bottom panel all galaxies with precise distances are supplied with error bars indicating where the galaxy should be situated if their distance from the observer changes by $\pm 1\sigma_D$.